

CORAGEN[®]

INSECTICIDE

Coragen[®] insecticide quick reference guide.



Coragen[®] insecticide is powered by Rynaxypyr[®], a unique mode of action that provides remarkable plant protection that's unlike any other chemistry. It delivers reliable, consistent, long-lasting protection against key insects. As a Group 28 product, Coragen[®] insecticide brings the added benefit of having no cross-resistance to other chemistries.

Packaging: 6 L jug x 2/case

Rainfast: 1 hour

Provides Extended Control: 7 to 21 days (depending on rate and when weather and temperature are optimal)

Staging: Controls hatching insects all the way through to adult stages.

Tank-mixable: Very stable in the tank and very tank-mixable. Refer to product label for tank-mixing instructions, and to Tank-Mixing Policy for Supported Tank-Mixes

Classified as a Reduced Risk Product: Has minimal impact on beneficial insects and pollinators when applied as directed by the product label¹.

Temperature: Can be applied in a wide range of temperatures (between 4°C and 40°C)

Water Volume: 10 Gal/ac

Crops	PHI	Insects
Alfalfa, sweet clover	1 day	Alfalfa weevil (suppression only)
Oilseeds: Canola, Mustard, Sunflower, Safflower	1 day	Bertha armyworm, diamondback moth, cutworms, swede midge, imported cabbageworms, sunflower head moth
Pulses: Chickpeas, Lentils, Peas, Beans	1 day	Armyworms, cutworms, grasshoppers, cabbage loopers, corn earworms, European corn borers, banded sunflower moth
Cereals	1 day	Grasshoppers, cutworms, armyworms, corn earworm, European corn borer
Sweet Corn	1 day	Armyworm, fall armyworm, variegated cutworm, corn earworm, Western bean cutworms, European corn borer
Seed Corn	1 day	Black cutworm
Field Corn, Popcorn	14 days	Black cutworm
Flax	1 day	Bertha armyworm, cutworms
Grass Forage, Hay	0 days	Bertha armyworm, cutworms
Potatoes	1 day	Colorado potato beetle

Questions? Ask your retailer about Coragen[®] insecticide today. For more information, call 1-833-362-7722 or visit FMCcrop.ca

¹In line with Integrated Pest Management and Good Agricultural Practices, insecticide applications should be made when pollinators are not foraging to avoid unnecessary exposure.

Always read and follow label directions.
Member of CropLife Canada.

FMC, Coragen and Rynaxypyr are trademarks of FMC Corporation. © 2018 FMC Corporation.